



**Missouri State**<sup>™</sup>  
U N I V E R S I T Y

**BearWorks**

---

Articles by College of Agriculture Faculty

Darr College of Agriculture

---

2009

## Captive Elephant Medicine: Recent Development

Ellen Wiedner

Dennis L. Schmitt  
*Missouri State University*

Follow this and additional works at: <https://bearworks.missouristate.edu/articles-coa>

---

### Recommended Citation

Wiedner, E., and D. L. Schmitt. "Captive elephant medicine: recent developments." *Gajah* 31 (2009): 25-28.

This article or document was made available through BearWorks, the institutional repository of Missouri State University. The work contained in it may be protected by copyright and require permission of the copyright holder for reuse or redistribution.

For more information, please contact [BearWorks@library.missouristate.edu](mailto: BearWorks@library.missouristate.edu).

## Captive Elephant Medicine: Recent Developments

Ellen Wiedner<sup>1</sup> and Dennis L. Schmitt<sup>1,2</sup>

<sup>1</sup>*Ringling Bros. and Barnum & Bailey Center for Elephant Conservation, Polk City, USA*

<sup>2</sup>*Agriculture Department, Missouri State University, Springfield, USA*

Several new challenges in elephant medicine have emerged over the past several years. Arguably the most critical is elephant endotheliotropic herpesvirus (EEHV), now recognized as a significant threat to captive Asian elephant populations in range countries, Europe and Asia (Cracknell *et al.* 2007; Reid *et al.* 2007; Richman 2007). Characterized by high mortality as well as stillbirth and abortion, the disease primarily affects juveniles between the ages of 1 and 4 years. Affected animals typically present with mild signs of lethargy which progress within hours or days to cranial and respiratory edema, facial cyanosis, and ultimately death, caused by myocardial failure and damage to the endothelial lining of the microvasculature leading to severe internal hemorrhage. Diagnosis is achieved via PCR testing of whole blood samples from actively viremic animals (Richman *et al.* 1999; Richman 2007). Current treatment recommendations include large doses of the antiviral drug famcyclovir, along with supportive care (Schmitt *et al.* 2000). However, in several cases, elephants have died despite early and aggressive treatment for reasons that remain unclear.

EEHV is a member of the genus Proboscivirus, and little is known about its epidemiology, modes of transmission, or patterns of shedding and latency. Three distinct strains were initially classified: EEHV 1a, found in both Asian and African elephants, EEHV 1b, which is apparently specific to Asian elephants, and EEHV 2, documented both in healthy wild African elephants and in two African elephants in North America who succumbed to hemorrhagic disease. Recently, however, several new strains of EEHV have been recognized, of which some are associated with disease and others are not (Garner *et al.* 2009). The relationships between the new strains are under investigation, and it is still unclear whether they are infective and/or pathologic to

both elephant species.

Efforts to establish a databank of elephant serum from North American animals are being performed by several institutions in conjunction with the EEHV laboratory at the National Zoo in Washington, DC in order to identify epidemiological patterns of transmission, seroconversion, and shedding. A recently developed ELISA test from the National Zoo against EEHV1a antibodies allows identification of at least some of animals that have been exposed to the disease. Current estimates indicate that more than 10% of North American elephants are seropositive for EEHV suggesting that seroconversion occurs with subclinical disease (Latimer 2007). It is hoped that measurement of viral titers will eventually help identify animals most likely to shed disease -- information that could be used in managing the movement of captive animals between institutions. A similar databank documenting EEHV titers elephants in Europe and in range countries is planned (Cracknell *et al.* 2007; Reid *et al.* 2006).

Another area of intense research in elephant medicine is tuberculosis (*Mycobacterium tuberculosis*), due, in part to the growing need for better and more reliable methods of diagnosis. Trunk washes, while 100% specific and still the gold standard for diagnosis, have less than ideal sensitivity because elephants only shed mycobacterial organisms intermittently. Diagnostic techniques useful in other species including chest radiographs, skin testing, and interferon gamma levels are either not feasible or not reliable in elephants. (Miller 2007) Although several new serological tests have appeared on the market, their validity remains unproven, and the possibility of false positives using these tests in animals exposed to atypical mycobacteria or even non-mycobacterial cross-reacting antigens

is concerning. (Cousins & Florisson 2005) Because treatment of a positive animal is a significant endeavor, and because legislation in several European countries requires euthanasia of any zoo animal with even a suspicion of tuberculosis (*M. tb*), false positives have grave consequences.

Treatment of *M. tb* in elephants still continues to be a conundrum as well. In the United States, treatment requirements are based on older guidelines developed for human with *M. tb*, and evidence now exists indicating that several of these drugs when dosed to produce human serum levels, cause severe toxicity in elephants. The options of utilizing alternative treatment regimes now standard in human *M. tb* therapy, including shorter duration of treatment, intermittent dosing schedules, and the use of other, less toxic drugs, are currently being investigated in elephants (Wiedner & Schmitt 2007).

With several demographic models forecasting the inevitable loss of elephants in captivity due to aging populations and a low birth rate (Wiese and Willis 2004), ongoing research in assisted reproductive technologies has continued unabated (Hermes *et al.* 2007). Reproductive ultrasound of both male and female elephants, used to identify reproductive tract pathology, monitor developing pregnancies, and provide information about breeding soundness and infertility, has become an invaluable technique for institutions committed to breeding their animals (Hildebrandt *et al.* 1998, 2000, 2006).

Improvements in endocrine monitoring now permit tracking of the elephant's estrus cycle (Brown 2000). Elephants are unique in having two surges of luteinizing hormone (LH) during the follicular phase of the estrous cycle, the first is non-ovulatory and second LH surge induces ovulation. Newer assays for LH and progesterone enable veterinarians and elephant managers to accurately predict the occurrence of this second LH peak and facilitate better planning of both natural and artificial inseminations. (Dahl *et al.* 2004) A current limitation to using artificial insemination in Asian elephants is the difficulty of routinely freezing and storing semen for later

use, spurring research on sperm physiology, the use of extenders and freezing protocols.

Ironically, the development of an elephant contraceptive vaccine has also been an important recent development (Barber *et al.* 2001; Fayerer-Hoskins *et al.* 2000). Tests of a recently developed zona pellucida vaccine in Africa indicate that use of this immuno-contraceptive may be an alternative to permanent removal from the reproductive pool in areas where elephant populations need to be controlled (Barber *et al.* 2001; Fayerer-Hoskins *et al.* 2000).

A final area of active investigation is basic anatomy. Gross imaging techniques such as MRI and CT have enabled researchers to produce detailed anatomic maps of elephant joints, limbs, and muscle as well as document locomotion (Tsaopoulous 2008; Hakeem *et al.* 2008). Sophisticated camera recording techniques have unlocked clues about elephant gait and locomotion (Hutchinson *et al.* 2003, 2006; Ren & Hutchinson 2007). Microscopic techniques including histology and scanning electron microscopy have been used to analyze the placenta, the nervous system, and the cushions of the feet (Hoffman *et al.* 2005; Shoshani *et al.* 2006; Weissengruber *et al.* 2006; Witter *et al.* 2007). The potential clinical value of this type of information is high, since anatomic studies that delineate the ranges of normal help in both the recognition and understanding of pathology. As conservation of elephants takes on ever greater urgency, the ability of veterinarians to keep captive populations healthy becomes more important as well.

## References

- Barber, M.R., Lee, S.M., Steffens, W.L., Ard, M. & Fayerer-Hosken R.A. (2001) Immunolocalization of zona pellucida antigens in the ovarian follicle of dogs, cats, horses, and elephants. *Theriogenology* **55**: 1705-1717.
- Brown, J.L. (2000) Reproductive endocrine monitoring of elephants: An essential tool for assisting captive management. *Zoo Biology* **19**: 347-367.

- Cousins, D.V. & Florisson, N. (2005) A review of tests available for use in the diagnosis of tuberculosis in non-bovine species. *Review of Science and Technology* **24**: 1039-1059.
- Cracknell, J., Routh, A., Field, D., Schaftenaar, W., Fickel, J., Hildebrandt, T.B. & H.S. (2007) Elephant endotheliotropic herpesvirus (Eehv) in the United Kingdom: An update including the development of the National Eehv Management Programme. *Verhandlungsbericht Erkrankungen Zootiere* **43**.
- Dahl, N.J., Olson, D., Schmitt, D.L., Blasko, D.R., Kristipati, R. & Roser, J.F. (2004) Development of an enzyme-linked immunosorbent assay (ELISA) for luteinizing hormone (LH) in the elephant (*Loxodonta africana* and *Elephas maximus*). *Zoo Biology* **23**: 65-78.
- Fayrer-Hosken, R.A., Grobler, D., Van Altena, J.J., Bertschinger, H.J. & Kirkpatrick, J.F. (2000) Immunocontraception of African elephants. *Nature* **407**: 149.
- Garner, M.M., Helmick, K., Ochsenreiter, J., Richman, L.K., Latimer, E., Wise, A.G., Maes, R.K., Kiupel, M., Nordhausen, R.W., Zong, J.C. & Hayward, G.S. (2009) Clinico-pathologic features of fatal disease attributed to new variants of endotheliotropic herpesviruses in two Asian elephants (*Elephas maximus*). *Veterinary Pathology* **46**: 97-104.
- Hakeem, A.Y., Hof, P.R., Sherwood, C.C., Switzer III, R.C., Rasmussen, L.E.L. & Allman, J.M. (2005) Brain of the African elephant (*Loxodonta africana*): Neuroanatomy from magnetic resonance images. *The Anatomical Record* **287A**: 1117-1127.
- Hermes, R., Goeritz, F., Streich, W.J. & Hildebrandt T.B. (2007) Assisted reproduction in female rhinoceros and elephants - Current status and future perspective. *Reproduction of Domestic Animals* **42**: 33-44.
- Hildebrandt, T.B., Goeritz, F., Hermes, R., Reid, C., Denhard, M. & Brown, J.L. (2006) Aspects of the reproductive biology and breeding management of Asian and African elephants *Elephas maximus* and *Loxodonta africana*. *International Zoo Yearbook* **40**: 20-40.
- Hildebrandt, T.B., Goeritz, F., Pratt, N.C., Brown, J.L., Montali, R.J., Schmitt, D.L., Fritsch, G. & Hermes, R. (2000) Ultrasonography of the urogenital tract in elephants (*Loxodonta africana* and *Elephas maximus*): An important tool for assessing female reproductive function. *Zoo Biology* **19**: 321-332.
- Hildebrandt, T.B., Goeritz, F., Pratt, N.C., Schmitt, D.L., Quandt, S., Raath, J. & Hofmann, R.R. (1998) Reproductive assessment of male elephants (*Loxodonta africana* and *Elephas maximus*) by ultrasonography. *Journal of Zoo and Wildlife Medicine* **29**: 114-128.
- Hoffmann, A., Loderstedt, S., Eulenberger, K., Fluegger, M., Junhold, J. & Seeger, J. (2005) The epithelial layer of the term placenta of the Asian elephant (*Elephas maximus*): a light and scanning electron microscopic study. *Anatomia, Histologia and Embryologia* **34**: 19-20.
- Hutchinson, J.R., Famini, D., Lair, R. & Kram, R. (2003) Are fast-moving elephants really running? *Nature* **422**: 493-494.
- Hutchinson, J.R., Schwerda, D., Famini, D.J., Dale, R.H.I., Fischer, M.S. & Kram, R. (2006) The locomotor kinematics of Asian and African elephants: Changes with speed and size. *Journal of Experimental Biology* **209**: 3812-3827.
- Latimer, E., Richman, L., Garner, M., Helmick, K., Zong, J.-C., Jeaggans, S. & Hayward, G. (2007) Elephant endotheliotropic herpesviruses: update. In: *International Elephant Foundation Annual Meeting Abstracts*.
- Miller, M.A. (2007) Current Diagnostic methods for tuberculosis in zoo animals. In: Fowler, M.E. & Miller, R.E. (eds.) *Zoo and Wild Animal Medicine. Current Therapy. 6. ed.* Saunders. pp 10-19.
- Reid, C.E., Marx, N., Fickel, J., Goeritz, F., Hunt, M., Thy, N., Reynes, J.M., Schaftenaar, W. &

Hildebrandt, T.B. (2005) Endotheliotropic herpes in Asia: The impact on captive and wild Asian elephant populations. *Verhandlungsbericht Erkrankungen Zootiere* **42**: 273 (Abstract only).

Ren, L. & Hutchinson, J. (2007) Three-dimensional locomotor dynamics of African (*Loxodonta africana*) and Asian (*Elephas maximus*) elephants. *Comparative Biochemistry and Physiology A: Comparative Physiology* **146**: S107-S127.

Richman, L.K. (2007) Elephant herpesviruses. In: *Zoo and Wild Animal Medicine. Current Therapy, 6. ed.* Fowler, M.E. & Miller, R.E. (eds.). Saunders. pp. 349-354.

Richman, L.K., Montali, R.J., Gardner, J.M., Kennedy, M.A., Lehnhardt, J., Hildebrandt, T., Schmitt, D., Hardy, D., Alcendor, D.J. & Hayward, G.S. (1999) Novel endotheliotropic herpesviruses fatal for Asian and African elephants. *Science* **283**: 1171-1176.

Schmitt, D.L., Hardy, D.A., Montali, R.J., Richman, L.K., Lindsay, W.A., Isaza, R. & West, G. (2000) Use of famciclovir for the treatment of endotheliotropic herpesvirus infections in Asian elephants (*Elephas maximus*). *Journal of Zoo and Wildlife Medicine* **31**: 518-522.

Schmitt, D.L. & Hildebrandt, T.B. (1998) Manual collection and characterization of semen from Asian elephants (*Elephas maximus*). *Animal Reproduction Science* **53**: 309-314.

Shoshani, J., Kupsky, W.J. & Marchant, G.H. (2006) Elephant brain Part I: Gross morphology, functions, comparative anatomy, and evolution. *Brain Research Bulletin* **70**: 124-170.

Tsaopoulos, D., Miller, C.E., Ren, L. & Hutchinson, J. (2008) A 3d musculoskeletal computer model of elephant fore and hindlimbs. *Comparative Biochemistry and Physiology, Part A* **15**: 574-592.

Weissengruber, G.E., Egger, G.F., Hutchinson, J.R., Groenewald, H.B., Elsaesser, L., Famini, D. & Forstenpointner, G. (2006) The structure

of the cushions in the feet of African elephants (*Loxodonta africana*). *Journal of Anatomy* **209**: 781-792.

Weissengruber, G.E., Fuss, F.K., Egger, G., Stanek, G., Hittmair, K.M. & Forstenpointner, G. (2006) The elephant knee joint: morphological and biomechanical considerations. *Journal of Anatomy* **208**: 59-72.

Wiedner, E. & Schmitt, D.L. (2007) Preliminary report of side effects seen in elephants treated for tuberculosis (*Mycobacterium tuberculosis*). In: *Internat. Elephant Foundation Annual Meeting*.

Wiese, R.J. & Willis, K. (2004) Calculation of longevity and life expectancy in captive elephants. *Zoo Biology* **23**: 365-373.

Witter, K., Egger, G.F. & Boeck, P. (2007) Renaut bodies in nerves of the trunk of the African elephant, *Loxodonta africana*. *Journal of Morphology* **268**: 414-422.

Corresponding author's email:  
ewiedner@feldinc.com



Sundara with her mother  
Photo courtesy of Feld Entertainment