



---

College of Natural and Applied Sciences

---

2013

## **Inactivation of Tor proteins affects the dynamics of endocytic proteins in early stage of endocytosis**

B Tenay

*MSU Graduate Student*

E Kimberlin

*MSU Undergraduate*

M Williams

*MSU Undergraduate*

D Juliette

*MSU Undergraduate*

J Fakilahyel

*MSU Undergraduate*

*See next page for additional authors*

Follow this and additional works at: <https://bearworks.missouristate.edu/articles-cnas>

---

### **Recommended Citation**

Tenay, Brandon, Evin Kimberlin, Michelle Williams, Juliette Denise, Joshua Fakilahyel, and Kyoungtae Kim. "Inactivation of Tor proteins affects the dynamics of endocytic proteins in early stage of endocytosis." *Journal of biosciences* 38, no. 2 (2013): 351-361.

This article or document was made available through BearWorks, the institutional repository of Missouri State University. The work contained in it may be protected by copyright and require permission of the copyright holder for reuse or redistribution.

For more information, please contact [BearWorks@library.missouristate.edu](mailto: BearWorks@library.missouristate.edu).

---

**Authors**

B Tenay, E Kimberlin, M Williams, D Juliette, J Fakilahyel, and Kyoungtae Kim

---

# Inactivation of Tor proteins affects the dynamics of endocytic proteins in early stage of endocytosis

BRANDON TENAY, EVIN KIMBERLIN, MICHELLE WILLIAMS, JULIETTE DENISE, JOSHUA FAKILAHYEL  
and KYOUNGTAE KIM\*

*Department of Biology, Missouri State University, 901 South National Ave, Springfield,  
MO 65897, USA*

*\*Corresponding author (Fax, 417-836-5126; Email, [kkim@missouristate.edu](mailto:kkim@missouristate.edu))*

Tor2 is an activator of the Rom2/Rho1 pathway that regulates  $\alpha$ -factor internalization. Since the recruitment of endocytic proteins such as actin-binding proteins and the amphiphysins precedes the internalization of  $\alpha$ -factor, we hypothesized that loss of Tor function leads to an alteration in the dynamics of the endocytic proteins. We report here that endocytic proteins, Abp1 and Rvs167, are less recruited to endocytic sites not only in *tor2* but also *tor1* mutants. Furthermore, we found that the endocytic proteins Rvs167 and Sjl2 are completely mistargeted to the cytoplasm in *tor1* $\Delta$ *tor2*<sup>ts</sup> double mutant cells. We also demonstrate here that the efficiency of endocytic internalization or scission in all *tor* mutants was drastically decreased. In agreement with the Sjl2 mislocalization, we found that in *tor1* $\Delta$ *tor2*<sup>ts</sup> double mutant cells, as well as other *tor* mutant cells, the overall PIP<sub>2</sub> level was dramatically increased. Finally, the cell wall chitin content in *tor2*<sup>ts</sup> and *tor1* $\Delta$ *tor2*<sup>ts</sup> mutant cells was also significantly increased. Taken together, both functional Tor proteins, Tor1 and Tor2, are essentially required for proper endocytic protein dynamics at the early stage of endocytosis.

[Tenay B, Kimberlin E, Williams M, Denise J, Fakilahyel J and Kim K 2013 Inactivation of Tor proteins affects the dynamics of endocytic proteins in early stage of endocytosis. *J. Biosci.* **38** 351–361] DOI 10.1007/s12038-013-9326-7

---

## 1. Introduction

Target of rapamycin (TOR) proteins are found across eukaryotic organisms, including mammals and yeast. In mammals, it is known as mTOR (mechanistic TOR) and forms two complexes: mTOR complex 1 (mTORC1) and mTOR complex 2 (mTORC2) (Hall 2008; Laplante and Sabatini 2012). mTOR functions by modulating the activities of a diverse array of downstream effector proteins to control many specific signaling events within the cell (Kim *et al.* 2002; Loewith *et al.* 2002; Manning and Cantley 2007). In particular, it has been shown that mTORC1 is involved in controlling biosynthesis of proteins and growth (Laplante and Sabatini 2012), while mTORC2 plays a role in regulating cellular processes including cytoskeletal organization (Jacinto *et al.* 2004).

In *S. cerevisiae*, there are two known Tor proteins, Tor1 and Tor2, which were identified while screening mutants that showed resistance to rapamycin (Cafferkey *et al.* 1993). Tor proteins share a high homology (67%) at the nucleotide level (Helliwell *et al.* 1994). There are two distinct Tor complexes: TORC1 and TORC2. TORC1 is composed of Tor1 (or Tor2), Kog1 and Lst8, while TORC2 contains Tor2, Avo1, Avo2, Avo3, Bit61 and Lst8 (Loewith *et al.* 2002; Wedaman *et al.* 2003). Tor1 is a PIK-related protein kinase and, similar to its mammalian counterpart mTor, regulates the synthesis of proteins, which is required for the cell cycle progression from G1 to S phase (Barbet *et al.* 1996). Although it has been known that Tor1 and Tor2 share many biological functions, a unique feature of Tor2 similar to mTORC2 is that Tor2 regulates the reorganization of the actin cytoskeleton (Schmidt *et al.* 1996).

**Keywords.** Abp1; endocytosis; PIP<sub>2</sub>; scission; Tor2

Supplementary materials pertaining to this article are available on the *Journal of Biosciences* Website at <http://www.ias.ac.in/jbiosci/jun2013/supp/tenay.pdf>

Consistent with the functional divergence between the two kinases, their subcellular locations were found to be different from each other: Tor1 at the vacuole and Tor2 at the plasma membrane (Aronova *et al.* 2007; Berchtold and Walther 2009; Cardenas and Heitman 1995; Sturgill *et al.* 2008).

It is known that the actin cytoskeleton is required for the proper function of endocytosis in yeast (Kaksonen *et al.* 2005). Once an endocytic site has been selected, actin-independent endocytic proteins, including clathrin, are recruited first, which is followed by the recruitment of actin and actin-binding proteins, Abp1 and Arp2/3 complex (Kaksonen *et al.* 2003, 2005; Newpher *et al.* 2005). Abp1 and Arp2/3 complex stimulate actin polymerization, which facilitates the invagination of the plasma membrane (Kaksonen *et al.* 2005). Scission of the invagination takes place at the neck of the invagination with the help of Rvs161, Rvs167 and Vps1 (Idrissi *et al.* 2008; Kaksonen *et al.* 2005; Smaczynska-de *et al.* 2012). During and after scission of the endocytic vesicle, it appears that Sjl2 plays a role for the proper uncoating of endocytic proteins from the vesicle (Toret *et al.* 2008). With the known functional significance of the actin cytoskeleton in endocytosis, it has been suggested that the severity of actin defect correlates tightly with the severity of endocytic defect (Engqvist-Goldstein and Drubin 2003). For instance, cells containing mutant alleles of *TOR2* gene exhibited depolarization of the actin cytoskeleton and defects in their ability to internalize  $\alpha$ -factor pheromone via receptor-mediated endocytosis (deHart *et al.* 2003; Schmidt *et al.* 1996). However, the specific roles of Tor2 in the dynamics of endocytic proteins during early receptor-mediated endocytosis have not yet been documented. Recently, research has focused on quantitating the effects of gene mutations on the dynamics of endocytic proteins at the site of receptor-mediated endocytosis. In particular, with the help of real-time live-cell imaging of GFP-fusion proteins, researchers are able to quantitate the dynamic recruitment and internalization events of endocytic proteins, occurring at the endocytic sites. Using a similar approach we investigated effects of partial and complete loss of Tor function on the dynamics of Abp1-GFP, Rvs167-GFP and Sjl2-GFP, and here present that intact Tor proteins are required both for proper dynamics of these endocytic proteins and efficient internalization event.

## 2. Materials and methods

### 2.1 Yeast strain construction and media

Yeast strains used in this study are listed in table 1. Strains expressing GFP-fused proteins were constructed by integrating respective GFP sequence at the 3' end of the gene of interest as described previously (Longtine *et al.* 1998; Kim *et al.* 2006; Nannapaneni *et al.* 2010). Mutant strains *tor1* $\Delta$  in

which *TOR1* locus is replaced by a *LEU2* cassette (Helliwell *et al.* 1994), *tor2<sup>ts</sup>* (a Tor2 PI kinase temperature-sensitive mutant harbouring the *tor2<sup>kin-</sup>* allele) (Schmidt *et al.* 1996), *tor1* $\Delta$ *tor2<sup>ts</sup>* and Tor2 overexpression (Schmidt *et al.* 1996) were obtained from Michael Hall's lab. Since *TOR2* is an essential gene, Kunz *et al.* (1993) replaced the *TOR2* with a *ADE2* and introduced a plasmid with a mutation in the kinase domain of Tor2 to produce *tor2<sup>ts</sup>* strain. Tor2 overexpression strain (Schmidt *et al.* 1996) was constructed to overexpress Tor2 under the control of a GAL-promoter, and *tor2<sup>ts</sup>* cells overexpressing Tor2 were able to grow even at the restricted temperature. Tor2 overexpression strain was gift from Dr. Michael Hall (University of Basel). All yeast strains except Tor2 overexpressing cells were grown in yeast peptone dextrose (YPD) and cultured at either 30°C or 38°C. Tor2 overexpressing cells were grown in SD-URA-media-containing galactose (2%) in order to induce overexpression of Tor2. Cells expressing a PIP<sub>2</sub>-binding GFP-2XPH domain were constructed by introducing the GFP-2XPH PLC $\delta$  plasmid (Kindly provided by Scott Emr, Cornell University) by following one-step transformation protocol (Chen *et al.* 1992).

### 2.2 Spinning confocal microscopy

Time-lapse movies and still images of GFP labeled cells were made with a spinning disk confocal microscope. The system included an inverted Olympus 1X81 microscope, a Yokogawa CSUX1 spinning disk head, a 100 $\times$  numerical aperture (NA) 1.4 PlanApo oil objective, and an electron amplified CCD (ImageEM, Hamamatsu). The temperature at the stage was maintained at 30°C or 38°C. Images were focused at an equatorial plane of the cells. Time-lapse movies of GFP labeled proteins were captured at 2 frames/s for movie duration time of 1 min. PIP<sub>2</sub> was labeled by expressing GFP-2xPH (PLC $\delta$ ) as previously described (Stefan *et al.* 2002). To quantitate PIP<sub>2</sub> expression levels, 13 small-budded cells from each strain expressing GFP-2xPH (PLC $\delta$ ) were randomly chosen to determine the average total GFP fluorescence per cell. To visualize yeast cell wall, cells were stained with calcofluor as previously described (Bickle *et al.* 1998).

### 2.3 Measurement of patch lifetime at the membrane and kymograph

Patch lifetime at the membrane was determined by calculating the time spent by a GFP-fused protein at the membrane (from the time the patch appeared to the time at which the patch was internalized from its origin or disappeared), as previously described (Murphy *et al.* 2011). Mean patch lifespan of GFP-fused proteins at the membrane was determined by averaging lifespan of the florescences from more

**Table 1.** Yeast strains used for this study

Strain	Genotype	Source
KKY 917	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa tor1::LEU<sub>2-4</sub></i>	(Schmidt <i>et al.</i> 1996)
KKY 918	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> /YCplac<sub>111::tor2-21</sub><sup>ts</sup></i>	(Schmidt <i>et al.</i> 1996)
KKY 919	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor1::HIS<sub>3-3</sub> tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup></i>	(Schmidt <i>et al.</i> 1996)
KKY 920	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> /pJK5 (pSEYC68galp::TOR2)</i>	(Schmidt <i>et al.</i> 1996)
KKY 0051	<i>Mata his3 leu2 ura3 , ABP1-GFP-HIS3</i>	Invitrogen
KKY 0960	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa tor1::LEU<sub>2-4</sub>, ABP1-GFP-TRP1</i>	This study
KKY 0961	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> /YCplac<sub>111::tor2-21</sub><sup>ts</sup>, ABP1-GFP-TRP1</i>	This study
KKY 0962	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor1::HIS<sub>3-3</sub> tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup>, ABP1-GFP-TRP1</i>	This study
KKY 0963	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> /pJK5 (pSEYC68galp::TOR2), ABP1-GFP-TRP1</i>	This study
KKY 0661	<i>Mata his3 leu2 lys2 ura3, RVS167-GFP-HIS</i>	This study
KKY 1009	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa tor1::LEU<sub>2-4</sub>, RVS167-GFP-TRP1</i>	This study
KKY 1010	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> /YCplac<sub>111::tor2-21</sub><sup>ts</sup>, RVS167-GFP-TRP1</i>	This study
KKY 1011	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor1::HIS<sub>3-3</sub> tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup>, RVS167-GFP-TRP1</i>	This study
KKY 455	<i>Mata his3 leu2 met15 ura3, SJL2-GFP-HisMx6</i>	This study
KKY 1014	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor1::HIS<sub>3-3</sub> tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup>, SJL2-GFP-TRP1</i>	This study
KKY 1075	<i>Mata his3 ura3 leu2 trp1 lys2 pRS426GFP-2 PH(PLC<math>\delta</math>) (URA3)</i>	This study
KKY 1077	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa tor1::LEU<sub>2-4</sub>, pRS426GFP-2 PH(PLC<math>\delta</math>) (URA3)</i>	This study
KKY 1079	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup>, pRS426GFP-2 PH(PLC<math>\delta</math>) (URA3)</i>	This study
KKY 1081	<i>Mata leu2-3,112 ura3-52 rme1 trp1 his4 GAL+ HMLa ade2 tor1::HIS<sub>3-3</sub> tor2::ADE<sub>2-3</sub> / YCplac<sub>111::tor2-21</sub><sup>ts</sup>, pRS426GFP-2 PH(PLC<math>\delta</math>) (URA3)</i>	This study


than at least 20 patches. Patches that at any time appeared too close to another patch were excluded from our analysis. In order to directly visualize the duration and fluorescence intensity of the patch on the membrane, a kymographic representation of GFP-fused proteins in a single patch over time was made using Slidebook (v.5). Average fluorescence intensity of the GFP-patches was determined using Image J (v.1.4).

### 3. Results

#### 3.1 Partial and complete loss of Tor function affects Abp1 dynamics

It has been previously described that the rate of  $\alpha$ -factor internalization by receptor-mediated endocytosis (RME) was attenuated in *tor2*<sup>G2128R</sup> kinase mutant cells that express arginine instead of glycine at the 2128th residue within the Tor2 protein (deHart *et al.* 2003). In order to investigate how Tor2 plays a role in both the recruitment dynamics of endocytic proteins and the scission efficiency during RME,

a well-characterized RME marker Abp1 was fused with GFP and the dynamic behavior of Abp1-GFP in cells was analysed at both 30°C and 38°C, after recording time-lapse fluorescence images. Actin-binding protein Abp1 arrives at endocytic sites approximately 5–8 s before scission event, moves in a directed manner away from the endocytic site, and dissociates from the post-internalized endocytic patch to be reused at future endocytic sites (Kim *et al.* 2006; Toret *et al.* 2008). Representative fluorescent images of Abp1-GFP patches in Wt and *tor* mutant cells (*tor1*  $\Delta$ , *tor2*<sup>ts</sup>, *tor1*  $\Delta$  *tor2*<sup>ts</sup> and *tor2* overexpressing) are shown in figure 1A. We found that the mean membrane lifespans of Abp1-GFP in all *tor* mutant cells except Tor2 over expressing cells were similar to that of Wt cells at both 30°C and 38°C (8.2 $\pm$ 2.2 s) (figure 1B and C). These results suggest that Tor1 and functional Tor2-kinase domain are not required for both the targeting of Abp1-GFP to endocytic sites and its subsequent maturation event. However, overexpression of Tor2 led to a robust increase in Abp1-GFP lifespan (~30 s), more than 3-fold when compared with Wt, indicating that the overexpression negatively affects the maturation event of



**Figure 1.** Partial or complete inactivation of Tor proteins is accompanied by defects in endocytic internalization and relatively low amount of Abp1-GFP at endocytic sites. (A) Representative images of Abp1-GFP in Wt (KKY 0051), *tor1*Δ (KKY 0960), *tor2*<sup>ts</sup> (KKY 0961), *tor1*Δ*tor2*<sup>ts</sup> (KKY 0962), and Tor2 overexpression (KKY 0963) cells. (B and C) Mean membrane lifespan of Abp1-GFP at 30°C and 38°C. Abp1-GFP patches were analysed to determine the mean membrane lifespan as described in the methods section (n = number of patches). (D and E) Percent internalization or scission failure of Abp1-GFP patches at 30°C and 38°C. Patches were counted as internalized if they moved away from the site of recruitment. Patches that failed to show internalization were counted as unsuccessful patches in scission. Scission efficiency decreases in all tested *tor* mutant cells. (F) Kymograph representation of Abp1-GFP patch at the cell membrane over time in Wt and *tor1*Δ*tor2*<sup>ts</sup> cells. As shown in the kymograph, lifespans of both patches are similar, but the patch in *tor1*Δ*tor2*<sup>ts</sup> cell is dimmer. Fluorescence intensity of ten to twenty Abp1-GFP patches was analysed to get the mean fluorescence intensity. The mean fluorescence intensity of Abp1-GFP in Wt cells was normalized to 100%. An asterisk (\*) indicates a statistically significant difference ( $p < 0.05$ ) as compared to the mean fluorescence intensity of Wt Abp1-GFP patch.

Abp1-GFP. After recruited to endocytic sites, endocytic proteins that are physically associated with the endocytic vesicles are internalized. We, therefore, determined the extent to which Abp1-GFP patches undergo successful internalization or scission in *tor* mutant cells. Abp1-GFP patches that fail to move away from the site of recruitment were considered to be 'scission failure patches'. In most cases, Abp1-GFP appears to be dissociated from patches that fail to internalize, thereby those can also be considered as aborted patches. Thus, we concluded that the low rate of scission success in the *tor* mutants is attributed to abortion in endocytosis that might be caused by premature disassembly of Abp1. Strikingly, scission failure increased 4- to 9-fold at both 30°C and 38°C in *tor* mutant cells, with the highest failure rate in *tor1Δtor2<sup>ts</sup>* cells (~40%) (figure 1C and D). In addition to the low efficiency in endocytic scission, the fluorescence intensity of Abp1-GFP patches in *tor* mutants was dimmer than the patches observed in Wt cells, manifested by Kymographic representation shown in figure 1F and by a colocalization assay shown in supplementary figure 1A. We thus further quantitated average fluorescence intensity of Abp1-GFP per patch and found that the extent of Abp1-GFP fluorescence intensity in *tor* mutant cells decreased significantly (figure 1F). In particular, the fluorescence intensity of scission failure patches in *tor* mutant cells was far lower than that of a successfully internalizing Abp1-GFP patch in the same mutant cells (figure 1F). These results suggest that less Abp1-GFP is recruited to the endocytic sites in *tor* mutant cells than Wt. All together, we came to the conclusion that loss of Tor protein activities affects receptor-mediated endocytosis by significantly decreasing endocytic scission efficiency, which may be partially attributable to a suboptimal recruitment of Abp1 to the endocytic sites.


### 3.2 Impaired membrane targeting of Rvs167 in *tor* mutant cells

Upon observing the decrease in scission efficiency of Abp1-GFP in *tor* mutant cells both at 30 and 38°C, we set out to examine whether or not Tor proteins affect the dynamics of endocytic protein Rvs167, since Rvs167 has been shown to help membrane invagination and scission events (Kaksonen *et al.* 2005). Consistent with previous findings (Balguerie *et al.* 1999; Kaksonen *et al.* 2005), Rvs167-GFP in Wt cells was localized to cortical membrane patches. It was also observed that Rvs167-GFP, in *tor1Δ* and *tor2<sup>ts</sup>* mutant cells, is recruited to the cortical endocytic sites (figure 2A). However, it appears that the number of Rvs167-GFP patch recruited to the cortex was decreased (see below and supplementary figure 1B). However, surprisingly, we found that in *tor1Δtor2<sup>ts</sup>* mutant cells, Rvs167-GFP was completely mislocalized to the cytoplasm as punctate structures (figure 2A and supplementary

figure 1B). Often we observed patches that appeared associated to the cell cortex in the double mutant, but no patches were formed *de novo* at the cortex. The average membrane lifespans of Rvs167 in all *tor* mutant and Wt cells were similar to each other at 30°C (figure 2C). Although the membrane lifespans in *tor1Δ* and *tor2<sup>ts</sup>* cells were slightly decreased at 38°C, the time difference was not statistically significant. The extent to which Rvs167-GFP patches undergo successful internalization or scission in *tor* mutant cells was determined as described above. At both 30°C and 38°C, the averaged rate of scission failure was drastically increased both in *tor1Δ* and *tor2<sup>ts</sup>* cells (figure 2D and E), 2-fold (~20% at 30°C) and 4-fold (36% in *tor2<sup>ts</sup>* at 38°C), respectively, when compared to that of Wt (8.9%). However, the difference in scission rate between those cells was not statistically significant at 30°C, due to a high standard deviation in *tor1Δ* and *tor2<sup>ts</sup>*. The experimental results in figure 2E show that at 38°C the drastic difference in averaged scission rate between Wt and mutant strains is supported by a statistical analysis, as manifested by p-values being <0.05. Next, we quantified the number of Rvs167-GFP patches formed at the membrane cortex of a cell within a minute. Our analysis showed that the average numbers of Rvs167-GFP patches at the membrane both in *tor1Δ* and *tor2<sup>ts</sup>* cells were ~5, thus 25% of that of Wt (20±2.3 patches/min) (figure 2F and G). Kymograph representations of Rvs167-GFP patches in *tor* mutant cells showed a significant decrease in the fluorescence intensity of Rvs167-GFP (figure 2H). The quantification of Rvs167-GFP fluorescence intensity revealed that there was significantly less Rvs167-GFP being recruited to the endocytic sites in *tor* mutant cells when compared to Wt (figure 2I).

### 3.3 *Sjl2* mislocalization in *tor1Δtor2<sup>ts</sup>* mutant cells had a significant effect on membrane PIP<sub>2</sub> levels

Because Abp1 is known to be an important factor for the recruitment of Sjl2 to endocytic sites (Stefan *et al.* 2002, 2005), we examined Sjl2 localization in *tor* mutant cells that show defects in the proper dynamics of Abp1 and Rvs167 (figures 1 and 2). Sjl2 has been implicated in endocytosis via optimization of the plasma membrane PIP<sub>2</sub> levels (Stefan *et al.* 2002). As shown in figure 3, Sjl2-GFP was properly targeted to membrane cortex in Wt. In order to visualize Sjl2-GFP in *tor1Δ* and *tor2<sup>ts</sup>* cells, we introduced the GFP sequence at the 3' end of *SJL2*, but found that all colonies grown on selective media were found to be false-positives according to our analytical colony PCR (data not shown). Expression of Sjl2-GFP in *tor1Δtor2<sup>ts</sup>* mutant cells was successful, and in these, Sjl2-GFP was exclusively mistargeted to the cytoplasm, where it was seen as punctate patches (figure 3A and supplementary figure 1C). We also studied the localization of another synaptojanin-like protein,




**Figure 2.** Internalization defect of Rvs167-GFP and a significant reduction of Rvs167-GFP fluorescence intensity in *tor* mutant cells. (A) Representative images of Rvs167-GFP in Wt (KKY 0061), *tor1*Δ (KKY 1009), *tor2*<sup>ts</sup> (KKY 1010), and *tor1*Δ*tor2*<sup>ts</sup> (KKY 1011). (B and C) Mean membrane lifespan of Rvs167-GFP patches at 30°C and 38°C. 20 to 70 Rvs167-GFP patches were analysed to determine the mean membrane lifespan as described in the methods section. No significant defect in membrane lifespan was observed. (D and E) Percent scission failure of Rvs167-GFP patches at 30°C and 38°C. Scission efficiency decreases in all *tor* mutant cells. An asterisk (\*) indicates a statistically significant difference ( $p < 0.05$ ) as compared to the mean scission failure percentage of Wt Rvs167-GFP patch. (F and G) Mean number of Rvs167-GFP patches that form at the plasma membrane during 1 min movie duration. 20 to 70 Rvs167-GFP patches were analysed to determine the mean number of membrane patches that formed. In *tor* mutant cells, a significant reduction in number of Rvs167-GFP patches when compared to that in Wt (\* indicating  $p < 0.05$ ). (H) Kymograph representation of Rvs167-GFP patch at the cell membrane over time in Wt, *tor1*Δ, and *tor2*<sup>ts</sup>. (I) The mean fluorescence intensity of Rvs167-GFP patches ( $n = 20-70$ ). An asterisk (\*) indicates a statistically significant difference ( $p < 0.05$ ) as compared to the mean fluorescence intensity of Wt Rvs167-GFP patch.

Sjl1, in the *tor* double mutant, and found a similar abnormal Sjl1 localization in the cytoplasm (data not shown). It was shown that synaptojanins deficient cells (for example, *sjl1*Δ*sjl2*Δ double mutant) exhibit a robust increase in the

level of membrane PIP<sub>2</sub> (Stefan *et al.* 2002). Based on our finding of the severe mislocalization of Sjl1 and Sjl2 in the *tor* double mutant cells, we predicted that the double mutant cells would express an increased level of membrane PIP<sub>2</sub>,





**Figure 3.** Mistargeting of Sjl2 and hyperaccumulation of PIP<sub>2</sub> in *tor* mutant cells. (A) Representative images of Sjl2-GFP in Wt (KKY 455) and *tor1Δtor2<sup>ts</sup>* (KKY 1014) mutant cells. Noticeable cytoplasmic mistargeting of Sjl2-GFP in *tor1Δtor2<sup>ts</sup>* was observed. (B) Representative images of membrane PIP<sub>2</sub>. In order to visualize PIP<sub>2</sub>, the GFP-2xPH (PLCδ) plasmid was expressed in Wt (KKY 1075), *tor1Δ* (KKY 1077), *tor2<sup>ts</sup>* (KKY 1079), and *tor1Δtor2<sup>ts</sup>* (KKY 1081) cells. (C) The fluorescence intensity of PH-GFP for each bar was quantified in 13 randomly selected cells. The mean fluorescence intensity of PH-GFP in Wt cells was normalized to 100%. An asterisk (\*) indicates a statistically significant difference ( $p < 0.05$ ) as compared to the mean fluorescence intensity of Wt.

and found that this was the case; approximately a 11-fold increase in the fluorescence levels of GFP fused pleckstrin homology (PH) domain of PLCδ, a fluorescent reporter that binds PIP<sub>2</sub>, in the double mutant cells as shown in figure 3B and C. In addition, we found the average GFP fluorescence intensity from *tor1Δ* and *tor2<sup>ts</sup>* cells was significantly increased.

#### 3.4 Significant chitin accumulation in *tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>* mutant cells

Given that an endocytic blockade leads to an accumulation of cell wall chitin (Reyes *et al.* 2007), we examined cell wall chitin levels, using calcofluor white, in *tor* mutant cells. Representative images of calcofluor stained cell are shown in figure 4A. No change in the fluorescent intensity of calcofluor stained *tor1Δ* cells was observed when


compared to that of Wt cells, while chitin levels in *tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>* even in a permissive condition (at 30°C) were increased by 32% ( $p = 0.056$ ) and 94% ( $p < 0.0001$ ), respectively (figure 4B). In the non-permissive condition (at 38°C), both *tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>* mutant cells, as well as *tor1Δ* cells, exhibited significantly increased levels of chitin.

## 4. Discussion

Results from our study provide the first evidence that Tor proteins, Tor1 and Tor2, are implicated in the proper recruitment of certain endocytic proteins and the subsequent scission efficiency of endocytic vesicles based on the following observations. First, upon partial and complete loss of Tor activity, Abp1 appears to arrive at endocytic sites normally, but scission efficiency was decreased (figure 1D and E), which could be partly attributed to less recruitment of Abp1 to the sites (figure 1F). Second, we observed that in all *tor* mutants, including *tor1Δ*, *tor2<sup>ts</sup>*, and *tor1Δtor2<sup>ts</sup>* double mutants, amphiphysin Rvs167 recruitment to the endocytic site was affected, manifested by a drastic reduction (~75%) in the number of *de novo* endocytic site formations (figure 2F and G), as well as lower levels of Rvs167-GFP fluorescence intensity at the site (figure 2H and I). In addition, the phosphoinositide phosphatase Sjl2 was exclusively mistargeted to the cytoplasmic patch-like puncta in *tor1Δtor2<sup>ts</sup>* mutant cells (figure 3). Third, we concluded that the endocytic defects we observed are most likely due mainly to downregulation of Rom2 pathway via partial or complete Tor2 inactivation in our *tor2* mutant cells (*tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>*), which also leads to an increase in cell wall chitin level (figure 4).

#### 4.1 Less scission efficiency of endocytic vesicles carrying Abp1-GFP in *tor* mutants

Abp1, a late marker for receptor-mediated endocytosis, is recruited to endocytic sites after the early endocytic adaptor Sla1 has been recruited. Abp1-carrying endocytic vesicles are pinched off from the plasma membrane within 10 s after its recruitment. Then, Abp1 departs from the post-internalized vesicles for re-use (Kaksonen *et al.* 2003; Kukulski *et al.* 2012). Abp1 is known to be an activator of the Arp2/3 complex that alone weakly nucleates actin assembly (Goode *et al.* 2001). However, it has been postulated that Abp1 helps Arp2/3 complex to be recruited to the actin filaments, thereby promoting the nucleation activity of Arp2/3 complex (Quintero-Monzon *et al.* 2005). Unlike mammalian cells, where chemical disruption of actin assembly only partially inhibits endocytosis (Lunn *et al.* 2000; Moskowitz *et al.* 2003), actin polymerization in yeast has been shown to be essential for endocytosis, based on the fact that cells treated with Latrunculin A exhibited no endocytic invagination



**Figure 4.** *tor* mutant cells have increase amount of cell wall chitin. (A) Wt (KKY 0343), *tor1Δ* (KKY 0917), *tor2<sup>ts</sup>* (KKY 0918), *tor1Δtor2<sup>ts</sup>* (KKY 0919) cells were stained with calcofluor white to visualize chitin under the confocal microscopy at different temperatures. (B) The mean fluorescence intensity of calcofluor white per cell (n=10). An asterisk (\*) indicates statistically significant difference ( $p < 0.05$ ) between the calcofluor mean intensity in Wt and each *tor* mutant. A triple asterisk (\*\*\*) indicates statistically significant difference, where  $p < 0.0001$ .

(Kukulski *et al.* 2012). Interestingly, as shown in figure 1F, it was obvious that less Abp1 protein was found at endocytic sites in all *tor* mutant cells (*tor1Δ*, *tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>*). In particular, in those mutants we observed nearly 70–80% reduction in fluorescence intensity of Abp1-GFP at endocytic patches that showed endocytic failure, pointing to a correlation between the amount of Abp1 with the scission success of endocytic patches in those cells. Nevertheless, it is not plausible to assert that the observed defects are simply attributed to the reduced amount of Abp1 at the endocytic site due to the fact that cells deficient in Abp1 exhibited no severe endocytic defects (Galletta *et al.* 2008). Rather, in the case that the total amount of the Arp2/3 activator Abp1 is less than the threshold level, one can expect that other functionally redundant Arp2/3 activators compensate for the partial loss of Abp1, possibly inducing the final step of the endocytic event, namely, the pinching-off process. However, knowing that Abp1 physically interacts with many endocytic factors, including Sla1 (endocytic adaptor), Rvs167 (amphiphysin), and actin-regulating kinases Ark1 and Prk1 (Colwill *et al.* 1999; Cope *et al.* 1999; Lila and Drubin 1997; Warren *et al.* 2002), it is more reasonable to propose that Abp1 in general plays a significant role in serving as an organizer of efficient endocytic machinery at the endocytic site. In this view, it is highly likely that *tor* mutant cells exhibit

aberration in recruitment of key endocytic proteins (for example, Rvs167 and Sjl2, see below), which may result in building a functionally less competent endocytic machinery, and thereby conferring low scission success (figures 1 and 2). Given that, in addition to its overlapping function with Tor1, Tor2 is uniquely implicated in organization of the actin cytoskeleton (Schmidt *et al.* 1996), it is surprising to observe a similar scission defect in *tor1Δ* cells comparable to that of *tor2<sup>ts</sup>* cells. An aberration in actin organization in yeast is tightly correlated with endocytic defect, which is the case for *tor2<sup>ts</sup>* cells (Schmidt *et al.* 1996). Because no actin organization defect in *tor1* mutant cells has been reported (Schmidt *et al.* 1996), our explanation for the observed scission defect in the background of *tor1Δ* would be an indirect effect of the loss of *TOR1*. Nonetheless, the question of how a lesser amount of Abp1 is recruited to endocytic sites in *tor* mutant cells remains elusive. Unexpectedly we observed a significant increase in Abp1-GFP lifespan upon Tor2 overexpression (figure 1B and C). Our explanation for this phenotype would be that Tor2 overexpression causes a dominant negative effect on Abp1-GFP dynamics at the endocytic site. This effect could be attributed to one or both of the following conditions. Given Tor2 acts as a potent protein kinase, the overexpression of Tor2 may result in phosphorylation of endocytic factors non-specifically, which affects endocytic protein maturation. Alternatively, it may be that Tor2 interacts with endocytic factors non-specifically due to the excessive amount of Tor2, and thereby sequestering them not to be readily recruited to the endocytic site.

#### 4.2 Drastic reduction of Rvs167-GFP carrying endocytic sites in *tor* mutant cells

Yeast genome contains the amphiphysins homologs, Rvs161 and Rvs167, which form a heterodimer, playing an important role in endocytosis (Douglas *et al.* 2009; Ren *et al.* 2006). Endocytic function of Rvs167 resides in its BAR domain since expression of BAR domain alone in *rvs167Δ* cells leads to the restoration of functional endocytosis (Sivadon *et al.* 1997). Rvs167 is recruited to endocytic sites 2–3 s after Abp1 has been recruited (Kaksonen *et al.* 2005), and the targeting appears to be mediated via its BAR domain, which contain positive amino acid residues that bind to the anionic plasma membrane (Peter *et al.* 2004). Due to the fact that purified BAR domain proteins are able to induce membrane curvature or tubulation *in vitro* and that a significant fraction of endocytic patches in *rvs167Δ* cells exhibits a ‘retraction phenotype’, Rvs167 was thought to promote endocytic scission (Peter *et al.* 2004; Kaksonen *et al.* 2005; Takei *et al.* 1999). But this idea is currently being challenged by a recent report which suggests Rvs167 is not essentially required for endocytic scission based on electron tomography results that

most of invaginated endocytic pits undergo pinching-off process in the absence of Rvs167 (Kukulski *et al.* 2012). Rather, the study showed that Rvs167 is recruited to newly-formed shallow invaginations, determined to be ~50 nm deep in length and after its recruitment further invagination (up to 100 nm) occurs, concluding that Rvs167 functions in stimulating membrane tubulation. This important finding may provide a plausible explanation for our observations in this report; the inefficient recruitment of Rvs167-GFP (supplementary figure 1B) and reduced scission in *tor* mutant cells (figure 2). As stated earlier, Arp2/3 complex-mediated actin assembly stimulated by its activators including Abp1 at endocytic sites is essential for formation of invaginating pits. Due to a significant loss in the amount of Abp1 and a potential subsequent defect in forming functional endocytic machinery, we, to put it simply, speculate that the invaginated pit in *tor* mutant cells, especially in *tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>* cells, might be shorter (< 50nm) on average than in Wt cells. As a result, Rvs167 could not readily be recruited to those shallow invaginations of the plasma membrane, and in turn further induction of invagination may be ceased. The complete dislocation of Rvs167-GFP into the cytoplasm in the double mutant (*tor1Δtor2<sup>ts</sup>*) cells (figure 2 and supplementary figure 1B) may then arise from a condition where the vast majority of endocytic invaginations in the double mutant cells are too short in depth to be recognized by Rvs167.

#### 4.3 *Tor* mutants exhibited a drastic increase in PIP<sub>2</sub> levels and significant chitin accumulation

Sjl2-GFP, which normally localizes to endocytic sites, was completely mislocalized to the cytoplasm of the *tor* double mutant cells (figure 3 and supplementary figure 1C), as is the case for Rvs167-GFP. Sjl2 is a yeast synaptojanin-like proteins that hydrolyses phosphates of inositol ring of PIP<sub>2</sub> to form PI (phosphoinositide), and yeast genome contains two more Sjl2 homologs, Sjl1 and Sjl3 (Srinivasan *et al.* 1997; Stolz *et al.* 1998; Guo *et al.* 1999). Previously, a group of researchers reported that loss of Sjl2 leads to its cytoplasmic homolog Sjl1 to be recruited to endocytic sites to compensate for the loss of Sjl2, concluding that both proteins share the same enzymatic activity (Sun *et al.* 2007). In light of finding that the severe mislocalization of Sjl2 in the *tor* double mutant cells (figure 3), we investigated the localization of Sjl1 and found that the GFP-fused-Sjl1 is mainly located in the cytoplasm (data not shown). As such, we reasoned that membrane PIP<sub>2</sub> levels in the double mutant should increase due to both Sjl1 and Sjl2 not being detected at the membrane cortex. Indeed, according to our GFP-2XPH fluorescence intensity results (figure 3), there was a drastic elevation in PIP<sub>2</sub> level in all *tor* mutant cells (*tor1Δ*, *tor2<sup>ts</sup>*, and *tor1Δtor2<sup>ts</sup>*) compared to that in Wt cells.

Inappropriate balance in membrane PIP<sub>2</sub> levels can adversely affect endocytosis. Since many actin-binding endocytic proteins are known to physically interact with PIP<sub>2</sub> at endocytic sites, low levels of PIP<sub>2</sub> result in an actin assembly defect, accompanied by a significant blocking of endocytosis (Abe *et al.* 2008; Logan and Mandato 2006). High levels of PIP<sub>2</sub> are in general associated with hyperstimulation of actin assembly, which also causes endocytic deficiency (Sun *et al.* 2007). Intriguingly, the observed hyperaccumulation of PIP<sub>2</sub> appears not to be correlated with less recruitment of endocytic proteins (Abp1 and Rvs167) in our *tor* mutant cells (figures 1 and 2). We, at the moment, are not able to adequately explain the molecular basis of this discrepancy. However, given the fact that PIP<sub>2</sub> binds and activates the Rom2-Rho1 pathway that controls endocytosis, we speculate that abnormally high accumulation of PIP<sub>2</sub> in *tor* mutant cells may dysregulate the Rom2-mediated endocytosis. The yeast cell wall comprised mostly of chitin is implicated in protecting the cell from rapid changes in osmotic pressure and against environmental stresses. Upon sensing of cell wall weakening (or damage), yeast cells up-regulate chitin synthesis to stabilize the cell wall (Dallies *et al.* 1998). In conditions where the mutant Tor2 protein is partially and completely inactivated, we found that cell wall chitin levels are significantly increased up to 100% (figure 4). Given the fact that an endocytic blockade impairs internalization of Chs3, the chitin synthase subunit, leading to a significant increase in chitin synthesis (Reyes *et al.* 2007), it is plausible to postulate that the drastic increase in chitin levels in the *tor* mutant (*tor2<sup>ts</sup>* and *tor1Δtor2<sup>ts</sup>*) might be due to an incompetent endocytosis, at least, of Chs3.

#### Acknowledgements

We thank Dr Hall (University of Basel) for providing *tor* mutant strains (*tor1Δ*, *tor2<sup>ts</sup>*, *tor1Δtor2<sup>ts</sup>*, and *tor2* overexpressing mutant). This work was supported by a National Scientific Foundation Grant 0923024 (to K KIM) and by thesis funding from Missouri State University (B Tenay).

#### References

- Abe N, Inoue T, Galvez T, Klein L and Meyer T 2008 Dissecting the role of PtdIns(4,5)P<sub>2</sub> in endocytosis and recycling of the transferrin receptor. *J. Cell Sci.* **121** 1488–1494
- Aronova S, Wedaman K, Anderson S, Yates J, 3rd, and Powers T 2007 Probing the membrane environment of the TOR kinases reveals functional interactions between TORC1, actin, and membrane trafficking in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **18** 2779–2794
- Balguerie A, Sivadon P, Bonneau M and Aigle M 1999 Rvs167p, the budding yeast homolog of amphiphysin, colocalizes with actin patches. *J. Cell Sci.* **112** 2529–2537

- Barbet NC, Schneider U, Helliwell SB, Stansfield I, Tuite MF and Hall MN 1996 TOR controls translation initiation and early G1 progression in yeast. *Mol. Biol. Cell* **7** 25–42
- Berchtold D and Walther TC 2009 TORC2 plasma membrane localization is essential for cell viability and restricted to a distinct domain. *Mol. Biol. Cell* **20** 1565–1575
- Bickle M, Delley PA, Schmidt A and Hall MN 1998 Cell wall integrity modulates RHO1 activity via the exchange factor ROM2. *EMBO J.* **17** 2235–2245
- Cafferkey R, Young PR, McLaughlin MM, Bergsma DJ, Koltin Y, Sathé GM, Faucette L, Eng WK, Johnson RK and Livi GP 1993 Dominant missense mutations in a novel yeast protein related to mammalian phosphatidylinositol 3-kinase and VPS34 abrogate rapamycin cytotoxicity. *Mol. Cell. Biol.* **13** 6012–6023
- Cardenas ME and Heitman J 1995 FKBP12-rapamycin target TOR2 is a vacuolar protein with an associated phosphatidylinositol-4 kinase activity. *EMBO J.* **14** 5892–5907
- Chen DC, Yang BC and Kuo TT 1992 One-step transformation of yeast in stationary phase. *Curr. Genet.* **21** 83–84
- Colwill K, Field D, Moore L, Friesen J and Andrews B 1999 In vivo analysis of the domains of yeast Rvs167p suggests Rvs167p function is mediated through multiple protein interactions. *Genetics* **152** 881–893
- Cope MJ, Yang S, Shang C and Drubin DG 1999 Novel protein kinases Ark1p and Prk1p associate with and regulate the cortical actin cytoskeleton in budding yeast. *J. Cell Biol.* **144** 1203–1218
- Dallies N, Francois J and Paquet V 1998 A new method for quantitative determination of polysaccharides in the yeast cell wall. Application to the cell wall defective mutants of *Saccharomyces cerevisiae*. *Yeast* **14** 1297–1306
- deHart AK, Schnell JD, Allen DA, Tsai JY and Hicke L 2003 Receptor internalization in yeast requires the Tor2-Rho1 signaling pathway. *Mol. Biol. Cell* **14** 4676–4684
- Douglas LM, Martin SW and Konopka JB 2009 BAR domain proteins Rvs161 and Rvs167 contribute to *Candida albicans* endocytosis, morphogenesis, and virulence. *Infect. Immun.* **77** 4150–4160
- Engqvist-Goldstein AE and Drubin DG 2003 Actin assembly and endocytosis: from yeast to mammals. *Annu. Rev. Cell Dev. Biol.* **19** 287–332
- Galletta BJ, Chuang DY and Cooper JA 2008 Distinct roles for Arp2/3 regulators in actin assembly and endocytosis. *PLoS Biol.* **6**e1
- Goode BL, Rodal AA, Barnes G and Drubin DG 2001 Activation of the Arp2/3 complex by the actin filament binding protein Abp1p. *J. Cell Biol.* **153** 627–634
- Guo S, Stolz LE, Lemrow SM and York JD 1999 SAC1-like domains of yeast SAC1, INP52, and INP53 and of human synaptojanin encode polyphosphoinositide phosphatases. *J. Biol. Chem.* **274** 12990–12995
- Hall MN 2008 mTOR-what does it do? *Transplant. Proc.* **40** S5–S8
- Helliwell SB, Wagner P, Kunz J, Deuter-Reinhard M, Henriquez R and Hall MN 1994 TOR1 and TOR2 are structurally and functionally similar but not identical phosphatidylinositol kinase homologues in yeast. *Mol. Biol. Cell* **5** 105–118
- Idrissi FZ, Grottsch H, Fernandez-Golbano IM, Presciatto-Baschong C, Riezman H and Geli MI 2008 Distinct acto/myosin-I structures associate with endocytic profiles at the plasma membrane. *J. Cell Biol.* **180** 1219–1232
- Jacinto E, Loewith R, Schmidt A, Lin S, Ruegg MA, Hall A and Hall MN 2004 Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. *Nat. Cell Biol.* **6** 1122–1128
- Kaksonen M, Sun Y and Drubin DG 2003 A pathway for association of receptors, adaptors, and actin during endocytic internalization. *Cell* **115** 475–487
- Kaksonen M, Toret CP and Drubin DG 2005 A modular design for the clathrin- and actin-mediated endocytosis machinery. *Cell* **123** 305–320
- Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P and Sabatini DM 2002 mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* **110** 163–175
- Kim K, Galletta BJ, Schmidt KO, Chang FS, Blumer KJ and Cooper JA 2006 Actin-based motility during endocytosis in budding yeast. *Mol. Biol. Cell* **17** 1354–1363
- Kukulski W, Schorb M, Kaksonen M and Briggs JA 2012 Plasma membrane reshaping during endocytosis is revealed by time-resolved electron tomography. *Cell* **150** 508–520
- Kunz J, Henriquez R, Schneider U, Deuter-Reinhard M, Movva NR and Hall MN 1993 Target of rapamycin in yeast, TOR2, is an essential phosphatidylinositol kinase homolog required for G1 progression. *Cell* **73** 585–596
- Laplante M and Sabatini DM 2012 mTOR signaling in growth control and disease. *Cell* **149** 274–293
- Lila T and Drubin DG 1997 Evidence for physical and functional interactions among two *Saccharomyces cerevisiae* SH3 domain proteins, an adenylyl cyclase-associated protein and the actin cytoskeleton. *Mol. Biol. Cell* **8** 367–385
- Loewith R, Jacinto E, Wullschlegel S, Lorberg A, Crespo JL, Bonenfant D, Oppliger W, Jenoe P and Hall MN 2002 Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. *Mol. Cell* **10** 457–468
- Logan MR and Mandato CA 2006 Regulation of the actin cytoskeleton by PIP2 in cytokinesis. *Biol. Cell* **98** 377–388
- Longtine MS, McKenzie A, 3rd, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P and Pringle JR 1998 Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* **14** 953–961
- Lunn JA, Wong H, Rozengurt E and Walsh JH 2000 Requirement of cortical actin organization for bombesin, endothelin, and EGF receptor internalization. *Am. J. Physiol. Cell Physiol.* **279** C2019–C2027
- Manning BD and Cantley LC 2007 AKT/PKB signaling: navigating downstream. *Cell* **129** 1261–1274
- Moskowitz HS, Heuser J, McGraw TE and Ryan TA 2003 Targeted chemical disruption of clathrin function in living cells. *Mol. Biol. Cell* **14** 4437–4447
- Murphy ER, Boxberger J, Colvin R, Lee SJ, Zahn G, Loor F and Kim K 2011 Pil1, an eisosome organizer, plays an important role in the recruitment of synaptojanins and amphiphysins to facilitate receptor-mediated endocytosis in yeast. *Eur. J. Cell Biol.* **90** 825–833
- Nannapaneni S, Wang D, Jain S, Schroeder B, Highfill C, Reustle L, Pittsley D, Maysent A, Moulder S, McDowell R, et al. 2010

- The yeast dynamin-like protein Vps1: vps1 mutations perturb the internalization and the motility of endocytic vesicles and endosomes via disorganization of the actin cytoskeleton. *Eur. J. Cell Biol.* **89** 499–508
- Newpher TM, Smith RP, Lemmon V and Lemmon SK 2005 In vivo dynamics of clathrin and its adaptor-dependent recruitment to the actin-based endocytic machinery in yeast. *Dev. Cell* **9** 87–98
- Peter BJ, Kent HM, Mills IG, Vallis Y, Butler PJ, Evans PR and McMahon HT 2004 BAR domains as sensors of membrane curvature: the amphiphysin BAR structure. *Science* **303** 495–499
- Quintero-Monzon O, Rodal AA, Strokopytov B, Almo SC and Goode BL 2005 Structural and functional dissection of the Abp1 ADFH actin-binding domain reveals versatile in vivo adapter functions. *Mol. Biol. Cell* **16** 3128–3139
- Ren G, Vajjhala P, Lee JS, Winsor B and Munn AL 2006 The BAR domain proteins: molding membranes in fission, fusion, and phagy. *Microbiol. Mol. Biol. Rev.* **70** 37–120
- Reyes A, Sanz M, Duran A and Roncero C 2007 Chitin synthase III requires Chs4p-dependent translocation of Chs3p into the plasma membrane. *J. Cell Sci.* **120** 1998–2009
- Schmidt A, Kunz J and Hall MN 1996 TOR2 is required for organization of the actin cytoskeleton in yeast. *Proc. Nat. Acad. Sci. USA* **93** 13780–13785
- Sivadon P, Crouzet M and Aigle M 1997 Functional assessment of the yeast Rvs161 and Rvs167 protein domains. *FEBS Lett.* **417** 21–27
- Smaczynska-de R II, Allwood EG, Mishra R, Booth WI, Aghamohammadzadeh S, Goldberg MW and Ayscough KR 2012 Yeast dynamin Vps1 and amphiphysin Rvs167 function together during endocytosis. *Traffic* **13** 317–328
- Srinivasan S, Seaman M, Nemoto Y, Daniell L, Suchy SF, Emr S, De Camilli P and Nussbaum R 1997 Disruption of three phosphatidylinositol-polyphosphate 5-phosphatase genes from *Saccharomyces cerevisiae* results in pleiotropic abnormalities of vacuole morphology, cell shape, and osmohomeostasis. *Eur. J. Cell Biol.* **74** 350–360
- Stefan CJ, Audhya A and Emr SD 2002 The yeast synaptojanin-like proteins control the cellular distribution of phosphatidylinositol (4,5)-bisphosphate. *Mol. Biol. Cell* **13** 542–557
- Stefan CJ, Padilla SM, Audhya A and Emr SD 2005 The phosphoinositide phosphatase Sjl2 is recruited to cortical actin patches in the control of vesicle formation and fission during endocytosis. *Mol. Cell Biol.* **25** 2910–2923
- Stolz LE, Huynh CV, Thorner J and York JD 1998 Identification and characterization of an essential family of inositol polyphosphate 5-phosphatases (INP51, INP52 and INP53 gene products) in the yeast *Saccharomyces cerevisiae*. *Genetics* **148** 1715–1729
- Sturgill TW, Cohen A, Diefenbacher M, Trautwein M, Martin DE and Hall MN 2008 TOR1 and TOR2 have distinct locations in live cells. *Eukaryotic Cell* **7** 1819–1830
- Sun Y, Carroll S, Kaksonen M, Toshima JY and Drubin DG 2007 PtdIns(4,5)P2 turnover is required for multiple stages during clathrin- and actin-dependent endocytic internalization. *J. Cell Biol.* **177** 355–367
- Takei K, Slepnev VI, Haucke V and De Camilli P 1999 Functional partnership between amphiphysin and dynamin in clathrin-mediated endocytosis. *Nat. Cell Biol.* **1** 33–39
- Toret CP, Lee L, Sekiya-Kawasaki M and Drubin DG 2008 Multiple pathways regulate endocytic coat disassembly in *Saccharomyces cerevisiae* for optimal downstream trafficking. *Traffic* **9** 848–859
- Warren DT, Andrews PD, Gourlay CW and Ayscough KR 2002 Sla1p couples the yeast endocytic machinery to proteins regulating actin dynamics. *J. Cell Sci.* **115** 1703–1715
- Wedaman KP, Reinke A, Anderson S, Yates J, 3rd, McCaffery JM and Powers T 2003 Tor kinases are in distinct membrane-associated protein complexes in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **14** 1204–1220

MS received 16 January 2013; accepted 15 March 2013

Corresponding editor: VEENA K PARNAIK