



---

College of Natural and Applied Sciences

---

8-23-2021

## **Editorial: Asteroseismology in the Kepler Era**

Anthony E. Lynas-Gray

Andrzej S. Baran  
*Missouri State University*

Karen Kinemuchi

Follow this and additional works at: <https://bearworks.missouristate.edu/articles-cnas>

---

### **Recommended Citation**

Lynas-Gray, Anthony E., Andrzej S. Baran, and Karen Kinemuchi. "Asteroseismology in the Kepler Era." *Frontiers in Astronomy and Space Sciences* (2021): 141.

This article or document was made available through BearWorks, the institutional repository of Missouri State University. The work contained in it may be protected by copyright and require permission of the copyright holder for reuse or redistribution.

For more information, please contact [BearWorks@library.missouristate.edu](mailto: BearWorks@library.missouristate.edu).



# Editorial: Asteroseismology in the Kepler Era

Anthony E. Lynas-Gray<sup>1,2,3</sup>, Andrzej S. Baran<sup>4,5,6\*</sup> and Karen Kinemuchi<sup>7</sup>

<sup>1</sup>University College London, London, United Kingdom, <sup>2</sup>University of Oxford, Oxford, United Kingdom, <sup>3</sup>University of the Western Cape, Bellville, South Africa, <sup>4</sup>ARDASTELLA Research Group, Institute of Physics, Pedagogical University of Kraków, Kraków, Poland, <sup>5</sup>Department of Physical Science, Embry-Riddle Aeronautical University, Daytona Beach, FL, United States, <sup>6</sup>Department of Physics, Astronomy, and Materials Science, Missouri State University, Springfield, MO, United States, <sup>7</sup>New Mexico State University, Las Cruces, NM, United States

**Keywords:** kepler spacecraft, pulsating stars, asteroseismology, space photometry, fourier technique

## Editorial on the Research Topic

### Asteroseismology in the Kepler Era

The Kepler spacecraft, launched by the National Aeronautics and Space Administration (NASA) on 2009 March 7th, proved to be an important pioneering astronomical survey instrument, making crucial discoveries during its primary mission over the ensuing 4 years. Looking longitudinally along the Local Spiral Arm in the general direction of Cygnus maximised the number of nearby (and therefore bright) stars that could be monitored almost continuously over the lifetime of the mission. A 0.95-m mirror with detectors sensitive to wavelengths in range 430–890 nm offered the best prospect of detecting earth-sized planets with orbital periods of about one-year, transiting across the disks of solar-type stars. While the search for planets orbiting nearby stars, and those in habitable zones in particular, persuaded NASA to support the Kepler mission, it would be a mistake to suppose that asteroseismic studies constituted a secondary scientific objective. Stellar pulsation generates light-curve features which could be of the same order as the flux lost from a solar-type star when an earth-sized planet in its habitable zone transits across its disk; it is critical that the two are not confused. Asteroseismology therefore plays a role in the search for planets beyond the Solar System. Furthermore, the existence of a planetary system is likely to affect evolution and internal structure of the host star, thereby determining its asteroseismic properties. Searches for planets orbiting nearby stars, and asteroseismology of these objects, are therefore inextricably linked.

The Editors are extremely grateful to the authors and reviewers who have contributed papers on asteroseismology with the Kepler spacecraft. While only a tiny fraction of the scientific returns are presented, the sample is understood to be representative and serve to inspire and support future space missions which will achieve even more than the impressive results secured by the Kepler mission.

In particular, we look forward to theoretical developments which allow the full potential of asteroseismic observations to be realised.

## AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Lynas-Gray, Baran and Kinemuchi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

## OPEN ACCESS

### Edited and reviewed by:

Scott William McIntosh,  
National Center for Atmospheric  
Research (UCAR), United States

### \*Correspondence:

Andrzej S. Baran  
andrzej.baran@up.krakow.pl

### Specialty section:

This article was submitted to  
Stellar and Solar Physics,  
a section of the journal  
Frontiers in Astronomy and Space  
Sciences

**Received:** 11 May 2021

**Accepted:** 11 August 2021

**Published:** 23 August 2021

### Citation:

Lynas-Gray AE, Baran AS and  
Kinemuchi K (2021) Editorial:  
Asteroseismology in the Kepler Era.  
Front. Astron. Space Sci. 8:708266.  
doi: 10.3389/fspas.2021.708266